

(19)



JAPANESE PATENT OFFICE

PATENT ABSTRACTS OF JAPAN

(11) Publication number: **59005956 A**

(43) Date of publication of application: **12.01.1984**

(51) Int. Cl. **G01N 31/22**
G01N 21/75

(21) Application number: **57114889**

(22) Date of filing: **01.07.1982**

(71) Applicant: **OSAKA SODA CO LTD**

(72) Inventor: **TOKUDA SHINGO**
ASADA SHIGEO
YAMAUCHI SHINJI

(54) **QUANTITATIVE MEASUREMENT OF MINUTE AMOUNT OF CALCIUM COMPONENT IN AQUEOUS SOLUTION**

(57) Abstract:

PURPOSE: To measure the concn. of a minute amount of calcium in good preciseness by an absorption photometry without receiving the influence of a magnesium salt or the like, by a method wherein glycine or aspartic acid is added and alizarin acid black SN is further added to perform colorimetric analysis.

CONSTITUTION: After a NaOH 10% solution is added to an aqueous 7% Na₂SO₄ solution containing a

minute amount of calcium and magnesium to adjust the pH thereof to 11.5, an aqueous 1% glycine solution is added to and mixed in the pH adjusted solution. In the next step, an aqueous 0.1% alizarin acid black SN solution is added thereto and pure water is added to measure absorbency by a spectrophotometer. When the aqueous solution of glycine or aspartic acid is added, the influence of a magnesium ion is not received even if it is present at pH of 11.5W13.0. The addition amount of glycine or aspartic acid is sufficient if added in an amount of 0.0001wt% or more. The addition amount of alizarin acid black SN as an indicator is 0.0001W0.01wt%.

COPYRIGHT: (C)1984,JPO&Japio

JP59005956A2: QUANTITATIVE MEASUREMENT OF MINUTE AMOUNT OF CALCIUM COMPONENT IN AQUEOUS SOLUTION

Country:	JP Japan
Kind:	A
Inventor:	TOKUDA SHINGO; ASADA SHIGEO; YAMAUCHI SHINJI;
Assignee:	OSAKA SODA CO LTD News, Profiles, Stocks and More about this company
Published/ Filed:	Jan. 12, 1984 / July 1, 1982
Application Number:	JP1982000114889
IPC Code:	G01N 31/22; G01N 21/75;
Priority Number:	July 1, 1982 JP1982000114889
Abstract:	<p>PURPOSE: To measure the concn. of a minute amount of calcium in good preciseness by an absorption photometry without receiving the influence of a magnesium salt or the like, by a method wherein glycine or aspartic acid is added and alizarin acid black SN is further added to perform colorimetric analysis.</p> <p>CONSTITUTION: After a NaOH 10% solution is added to an aqueous 7% Na₂SO₄ solution containing a minute amount of calcium and magnesium to adjust the pH thereof to 11.5, an aqueous 1% glycine solution is added to and mixed in the pH adjusted solution. In the next step, an aqueous 0.1% alizarin acid black SN solution is added thereto and pure water is added to measure absorbency by a spectrophotometer. When the aqueous solution of glycine or aspartic acid is added, the influence of a magnesium ion is not received even if it is present at pH of 11.5W13.0. The addition amount of glycine or aspartic acid is sufficient if added in an amount of 0.0001wt% or more. The addition amount of arizarin acid black SN as an indicator is 0.0001W0.01wt%.</p> <p>COPYRIGHT: (C)1984,JPO&Japio</p>
Family:	None
Other Abstract Info:	None



[View
Image](#)

1 page